

"I have to keep going, as there are always people on my track. I have to publish my present work as rapidly as possible in order to keep in the race. The best sprinters in this road of investigation are Becquerel and the Curies..." Letter to his mother (1902), Ernest Rutherford, New Zealand-English physicist (1871-1937)

"It was necessary at this point to find a new term to define this new property of matter manifested by the elements of uranium and thorium. I proposed the word radioactivity which has since become generally adopted; the radioactive elements have been called radio elements."

Marie Curie, Polish-French physicist (1867-1934)

"No-one really thought of fission before its discovery."

Lise Meitner, Austrian-Swedish physicist (1878-1968)

PHY100S (K. Strong) - Lecture 22 - Slide 1

Current Assignments ...

For today

- Read Sections 13.6-13.7, 14.1-14.2

For Lecture 23

- Sects. 14.3-14.4, 15.1-15.4, Ch. 17

Homework #4

- Posted March 7. **Late deadline 11 AM, March 29.**

Homework #5

- Posted March 21. Due 11:00 AM, Friday, April 5

Writing Assignment #2

- Posted Feb. 28. Due 11:00 AM, Thursday, April 4

Suggested Conceptual Exercises

- Chapter 14: 1,3,5,7,9,11,13,15,17,19,21,23,25,27,29

Tutorial #10

**Office hours:
3-4 Tuesdays
& Thursdays**

PHY100S (K. Strong) - Lecture 22 - Slide 2

Review of Lecture 21

Textbook, Sections 13.2-13.6

- The effect of observation
- Quantum nonlocality
- Quantum entanglement
- Toward a post-Newtonian worldview
- Spectroscopy and observing atomic spectra

PHY100S (K. Strong) - Lecture 22 - Slide 3

Plan for Lecture 22

Textbook, Sections 13.6-13.7

- Observing atomic spectra
- Models of the atom
- The quantum atom
- Energy transitions in atoms

Textbook, Sections 14.1-14.2

- The strong nuclear force
- Nuclear structure

PHY100S (K. Strong) - Lecture 22 - Slide 4

Examples of Spectra

The diagram shows three examples of spectra produced by a spectroscope. Each example shows a source of light on the left, a spectroscope in the middle, and the resulting spectrum on the right.

- Continuous spectrum:** A source of "White light" produces a "Continuous spectrum" (solid or liquid). The resulting spectrum is a continuous band of colors from violet to red.
- Line spectra:** A source of "Heated gas" produces "Bright lines characterizing the gas". The resulting spectrum shows discrete colored lines on a dark background.
- Dark absorption lines:** A source of "The same gas but cold transmitting light" produces "Dark absorption lines characterizing the gas". The resulting spectrum shows a continuous band of colors with dark lines superimposed.

Line spectra - dilute gas either emitting or absorbing

<http://www.amateur-spectroscopy.com/Spectroscope.htm>

PHY100S (K. Strong) - Lecture 22 - Slide 5

Interactive Examples of Spectra

- MiniSpectroscopy displays a visual representation (a "spectroscopy view") of a sample spectrum simultaneously with a graphical (intensity vs. wavelength) representation.

<http://mo-www.harvard.edu/Java/MiniSpectroscopy.html>



PHY100S (K. Strong) - Lecture 22 - Slide 6

Observing Atomic Spectra


- A gas can be excited by heat and by sending an electric current through it.
- But why does a thin gas emit only certain frequencies of light?

Textbook Figure 13.14

© 2010 Pearson Education, Inc.

PHY100S (K. Strong) - Lecture 22 - Slide 7

Models of the Atom




- To answer this question, we need to consider the atom again.
- Recall Lecture 8 - we discussed early models of the atom
 - billiard ball, plum pudding, planetary model
- 1913 - Bohr model of the atom (very briefly!)
 - stable orbits in which the electron can exist without radiating and thus not spiralling into the nucleus (but no theoretical justification)
 - each spectral line is due to energy lost when the electron falls from a higher to lower orbit (but only works for atoms with one electron)

PHY100S (K. Strong) - Lecture 22 - Slide 8

Standing Waves (towards a quantum model of the atom)

- Matter waves, like other waves, can form standing waves.
- There must be an integral number of half-waves for the wave to persist.
- The wave does not move along the string, but the string always has a wave pattern.




<http://faraday.physics.utoronto.ca/IYearLab/Intros/StandingWaves/Flash/sta2fix.html>

PHY100S (K. Strong) - Lecture 22 - Slide 9

The Quantum Atom

- The matter waves of the electron form circular standing waves.
- The wave must just fit around the atomic nucleus.
- The only allowed orbits of the electron are those for which this is true.



Copyright © 2007 Pearson Prentice Hall, Inc.

PHY100S (K. Strong) - Lecture 22 - Slide 10

Quantum States of the Hydrogen Atom: 10 possible electron distributions around the atom (there are more)

Textbook Figure 13.18

© 2010 Pearson Education, Inc.

These are 3D solutions to Schrodinger's Equation.

Darker shading indicates more intense matter field and higher probability that the electron will be there.

PHY100S (K. Strong) - Lecture 22 - Slide 11

Quantum States of Hydrogen Atom

- What does a hydrogen atom look like?
http://www.hydrogenlab.de/elektronium/HTML/einleitung_hauptseite_uk.html



- Hydrogen atom orbital viewer
<http://www.falstad.com/qmatom/>



Try these at home.

PHY100S (K. Strong) - Lecture 22 - Slide 12

Energy Level Diagrams

- Each quantum state is a standing wave with a specific frequency and a specific energy.
- The lowest energy level is called the ground state, and the rest are called excited states.

Textbook
Figures 13.18 & 13.19

© 2010 Pearson Education, Inc.

PHY100S (K. Strong) - Lecture 22 - Slide 13

Quantum Jumps

- When an electron makes a quantum jump from one quantum state to another, it makes a transition from one energy level to another.
- The atom emits/absorbs radiation if it jumps to a lower/higher energy level.
- Emitted/absorbed photon has energy


Textbook
Figure 13.20

© 2010 Pearson Education, Inc.
Emission of a photon

$$hf = E_{\text{high-energy state}} - E_{\text{low-energy state}}$$

PHY100S (K. Strong) - Lecture 22 - Slide 14

Absorption and Emission (schematic diagram of atom!)



http://superphysics.net/firms.com/pp_quantum_theory.html PHY100S (K. Strong) - Lecture 22 - Slide 15

All possible transitions between energy levels for the first five levels of the hydrogen atom

Energy levels

Textbook
Figures 13.21 & 13.22

Emitted photon frequencies

© 2010 Pearson Education, Inc.

PHY100S (K. Strong) - Lecture 22 - Slide 16

A certain type of atom has only four energy levels, as shown in the diagram. The "spectral lines" produced by this element are all visible, except for one ultra-violet line. The quantum jump that produces the UV line is

- (A) state 2 to 1 _____ 4
 (B) state 4 to 1 _____ 3
 (C) state 4 to 3 _____ 2
 (D) state 1 to 4 _____ 2
 (E) impossible to determine without further information _____ 1

PHY100S (K. Strong) - Lecture 22 - Slide 17

A certain type of atom has only four energy levels, as shown in the diagram. The total number of spectral lines emitted by this element is

- (A) 3 _____ 4
 (B) 4 _____ 3
 (C) 6 _____ 2
 (D) 10 _____ 2
 (E) impossible to determine without further information _____ 1

PHY100S (K. Strong) - Lecture 22 - Slide 18

Models of the Hydrogen Atom

Using spectroscopy to test models of the hydrogen atom

- <http://serc.carleton.edu/sp/compadre/interactive/examples/19268.html>

or

- <http://phet.colorado.edu/simulations/hydrogen-atom/hydrogen-atom.jnlp>



PHY100S (K. Strong) - Lecture 22 - Slide 19

A Simplified View of the Nucleus

Neutrons - uncharged

Protons - positively charged

Textbook
Figure 14.1

© 2010 Pearson Education, Inc.

Why doesn't electrical repulsion cause the nucleus to fly apart?

PHY100S (K. Strong) - Lecture 22 - Slide 20

The Strong Nuclear Force

- A "new" force, called the strong nuclear force, holds the nucleus together.
 - It acts over distances of about 10^{-15} m, attracting protons and neutrons.
- There is one other force, called the weak nuclear force, which is responsible for some forms of radioactive decay.
- Four fundamental forces – gravity, electromagnetism, strong and weak nuclear forces.
 - Responsible for the structure of our universe.
 - Every other force can be reduced to one of these four.

PHY100S (K. Strong) - Lecture 22 - Slide 21

The Size of the Strong Force

- The strong nuclear force is the strongest of the four forces. Why? quantum physics...
 - If a proton or neutron is confined to the nucleus, the Uncertainty Principle requires that its velocity be about 10% of the speed of light, so its kinetic energy is very large.
 - Similarly, if an electron is confined to the volume of an atom, its speed must be about 0.5% of the speed of light.
 - The energy difference between an electron moving at 0.5% of c and a proton moving at 10% of c yields the difference in strength between chemical and nuclear reactions.

PHY100S (K. Strong) - Lecture 22 - Slide 22

Some Definitions

- A nuclear reaction is any process that alters the structure of a nucleus.
 - Both protons and neutrons are important.
- Atomic number = the number of protons (also = number of orbital electrons)
 - This determines the element of the atom.
- The number of neutrons determines the isotope of the element.
- [Atomic] mass number = the total number of protons plus neutrons.
 - An isotope is labeled by its atomic number and its mass number.

PHY100S (K. Strong) - Lecture 22 - Slide 23

Periodic Table of Elements																	
Atomic number										Atomic mass ≈ mass number							
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Li	Be	B	C	N	O	F	Ne	Na	Mg	Al	Si	P	S	Cl	Ar	K	Ca
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
Fr	Ra	Ac	Rf	Sg	Bh	Hs	Mt	Ds	Rg	Cn	Fl	Mc	Lv	Ts	Og	Uu	Uub
For elements with no stable isotopes, the mass number of the isotope with the longest half-life is in parentheses.																	
Design and Interface Copyright © 1997 Michael Dayak (michael@dayak.com) http://www.ptable.com/																	
La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu			
Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr			

How do the mass and charge of a ^{14}C nucleus compare with those of a ^{12}C nucleus?

- (A) The mass of ^{14}C is 50% larger, while the charge is the same.
- (B) Both the mass and the charge of ^{14}C are the same as those of ^{12}C .
- (C) Both the mass and charge of ^{14}C are one-sixth larger.
- (D) The mass is the same, while the charge of ^{14}C is one-sixth larger.
- (E) The mass of ^{14}C is one-sixth larger, while the charge is the same.**

PHY100S (K. Strong) - Lecture 22 - Slide 25

How many protons and how many neutrons are in the $^{56}_{26}\text{Fe}$ (iron) nucleus?

- (A) 26 protons and 30 neutrons**
- (B) 26 protons and 56 neutrons
- (C) 30 protons and 26 neutrons
- (D) 56 protons and 26 neutrons
- (E) None of the above

$^{56}_{26}\text{Fe}$ indicates mass number Chemical symbol
atomic number = number of protons
mass number = number of protons + number of neutrons

PHY100S (K. Strong) - Lecture 22 - Slide 26